



This is a repository copy of *Compound heterozygous variants in IFT140 as a cause of nonsyndromic retinitis pigmentosa*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/123955/>

Version: Accepted Version

---

**Article:**

Low, T., Kostakis, A. and Balasubramanian, M. [orcid.org/0000-0003-1488-3695](https://orcid.org/0000-0003-1488-3695) (2017)  
Compound heterozygous variants in IFT140 as a cause of nonsyndromic retinitis pigmentosa. *Ophthalmic Genetics* . pp. 1-2. ISSN 1381-6810

<https://doi.org/10.1080/13816810.2017.1393827>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Compound heterozygous variants in IFT140 as a cause of non-syndromic**

### **Retinitis Pigmentosa**

Tisiana Low<sup>1</sup>, Anastassios Kostakis<sup>2</sup>, Meena Balasubramanian<sup>3</sup>

<sup>1</sup>Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK S5 7AU

<sup>2</sup>Department of Ophthalmology, Doncaster and Bassetlaw Hospital, Worksop, UK

<sup>3</sup>Sheffield Clinical Genetics Service, Sheffield Children's NHS Foundation Trust, Sheffield, UK S10 2TH

### **Correspondence:**

Dr Meena Balasubramanian; Sheffield Clinical Genetics Service; Sheffield Children's NHS Foundation Trust; Sheffield, UK; Ph: +44 114 2717025; Fax: +44 114 2737467; E-mail: meena.balasubramanian@nhs.net

### **Introduction**

Retinitis pigmentosa (RP) refers to a group of inherited disorders that affect the retina's ability to respond to light, leading to progressive visual loss. Retinitis pigmentosa sine pigmento is a variant of RP in which there is an absence of characteristic peripheral bone-spicule like pigmentary changes.

One of the genes found to be responsible for RP is IFT140, a ciliary transporter gene (OMIM \*614620). Homozygous and compound heterozygous IFT140 variants have commonly been reported in Mainzer-Saldino syndrome and Jeune syndrome.<sup>1</sup> However, in recent literature, non-syndromic IFT140-related RP have been reported.

IFT140 encodes a sub-unit of intraflagellar transport complex A (IFTA), which is involved in retrograde ciliary transport.<sup>2</sup> It was previously referred to as KIAA0590 and located on chromosome 16p13.3. It is highly expressed in kidneys with moderate expression in ovary, testis, lung, prostate. Schmidts et al., 2013 demonstrated high expression of Ift140 in renal and retinal tissue in mouse embryos.<sup>3</sup> Although the phenotype associated with IFT140 variants is still emerging, it appears to encompass

a variable spectrum ranging from non-syndromic, isolated RP (as demonstrated in this clinical report) to Short-rib thoracic dysplasia 9 with or without polydactyly (SRTD9; OMIM # 266920).

### **Case Report**

We present a case of a 22-year-old female who attended her opticians for a frequent headaches review. It was found that her visual fields were restricted, with a slightly abnormal retina with greying and mottling (Fig 1). However, night-time vision was not reduced. An optical coherence tomography suggested likely RP, and electrodiagnostic testing confirmed the diagnosis.

She was the first child of healthy, non-consanguineous, White European parents and had a younger sister who was fit and well with no family history of RP. A recent ophthalmology follow-up review found that the patient had visual acuities of 6/6 with patchy visual defects in bilateral eyes, but no significant decrease in visual fields.

### **Materials and Methods**

Genomic DNA was obtained from patient and both parents with appropriate consent. A total of 176 genes associated with retinal dystrophy were targeted using Retinal dystrophy v3 Agilent SureSelect Custom Design and sequenced on the HiSeq2500 (Illumina) system according to manufacturer's protocols. The target enrichment design consisted of coding region of the transcripts, including the immediate splice sites (+/- 5 bases) for these genes. Sequence data was subsequently mapped with GenomeAnalysis ToolKitLite-v2.0.39 (GATK) and with hg19 human genome as a reference. Known polymorphisms were filtered using bioinformatic analysis.

### **Results**

Compound heterozygous variants in IFT140 NM\_014714.3: c.4182G>C p.(Thr1394Thr) and c.212C>T p.(Pro71Leu) were identified, and confirmed by

Sanger sequencing. IFT140 c.4182G>C p.(Thr1394Thr) is a synonymous change which has not previously been reported in the literature or any of the mutation/ variant databases. However, it affects the last base of exon 30 and in silico prediction tools using Alamut Visual version 2.6 (Interactive Biosoftware, Rouen, France) indicates that this change may reduce or abolish the splice donor site of intron 30. IFT140 c.212C>T p.(Pro71Leu) has previously been reported in the compound heterozygous state in a patient with non-syndromic RP. The proline residue at codon 71 is highly conserved and in silico analysis using Polyphen2 and SIFT predicts this change to be damaging. Parental analyses confirmed that the father carried the IFT140 c.4182G>C p.(Thr1394Thr) and mother carried the IFT140 c.212C>T p.(Pro71Leu), showing that the variants are in trans in the proband.

## **Discussion**

IFT140 is a protein coding gene, variants in this gene are often associated with a group of autosomal recessive syndromic ciliopathies also known as short rib thoracic dysplasia 9 with or without polydactyly (SRTD9). Human IFT140 protein contains WD40 (WD) and TPR (tetratricopeptide) repeat domains. There does not appear to be specific genotype-phenotype correlation based on existing literature and variants within TPR and WD repeat domains in IFT140 protein give rise to both a syndromic ciliopathy and a non-syndromic RP. It has also been reported that the IFT140 missense variants in Mainzer-Saldino syndrome and Jeune syndrome were mainly found in the N-terminus and WD repeat domains.

The principle of genetics in RP is complex as locus and allelic heterogeneity exists. The use of multi-gene panels and next-generation sequencing currently yield an 80% success rate in the identification of the pathogenic variant(s). It is worth noting that individuals with RP often do not present with the classical appearance of bone spicule

pigmentation, which may lead to a delay in diagnosis. Indications such as poor night vision and positive family history should guide physicians to request electrodiagnostic tests for further investigation.

Since the use of next generation sequencing for retinal disorders, in clinical practice, we are able to identify more individuals with genetic forms of RP and provide diagnostic confirmation, information on prognosis and informed recurrence risks. This aids in genetic counselling, especially where there is no relevant family history and particularly helpful to clarify recurrence risks in unaffected family members. This report adds to the emerging literature of IFT140 causing a non-syndromic RP.

### **Figure Legend**

Fig 1. Retinal photographs demonstrating mottled appearance with greying of retina.

### **Acknowledgement**

We would like to thank the family for consenting to publication.

### **References**

- <sup>1</sup>Bifari IN, Elkhamary SM, Bolz HJ, et al. The ophthalmic phenotype of IFT140-related ciliopathy ranges from isolated to syndromic congenital retinal dystrophy. *Br J Ophthalmol*. 2015; **0**:1-5.
  - <sup>2</sup>Perrault I, Saunier S, Hanein S, et al. Mainzer-Saldino Syndrome Is a Ciliopathy Caused by IFT140 Mutations. *Am J Hum Genet*. 2012; **90**(5):864–870.
  - <sup>3</sup>Schmidts M, Frank V, Eisenberger T, et al. Combined NGS Approaches Identify Mutations in the Intraflagellar Transport Gene IFT140 in Skeletal Ciliopathies with Early Progressive Kidney Disease. *Hum Mutat*. 2013; **34**(5):714–724.
  - <sup>4</sup>Hull S, Owen N, Islam F, et al. Nonsyndromic retinal dystrophy due to bi-allelic mutations in the ciliary transport gene IFT140. *Invest Ophthalmol Vis Sci*. 2016; **57**:1053–1062.
  - <sup>5</sup>Neveling K, Collin R, Gilissen C, et al. Next-generation genetic testing for retinitis pigmentosa. *Hum Mutat*. 2012; **33**(6):963-972.
-